

LI-YORKE CHAOS FOR DENDRITE MAPS WITH ZERO TOPOLOGICAL ENTROPY AND ω -LIMIT SETS

GHASSEN ASKRI

ABSTRACT. Let X be a dendrite with set of endpoints $E(X)$ closed and let $f : X \rightarrow X$ be a continuous map with zero topological entropy. Let $P(f)$ be the set of periodic points of f . We prove that if L is an infinite ω -limit set of f then $L \cap P(f) \subset E(X)'$, where $E(X)'$ is the set of all accumulations points of $E(X)$. Furthermore, if $E(X)$ is countable and L is uncountable then $L \cap P(f) = \emptyset$. We also show that if $E(X)'$ is finite then any uncountable ω -limit set of f has a decomposition and as a consequence if f has a Li-Yorke pair (x, y) with $\omega_f(x)$ or $\omega_f(y)$ is uncountable then f is Li-Yorke chaotic.

1. Introduction

Let X be a compact metric space with metric d and $f : X \rightarrow X$ be a continuous map. Let $\mathbb{Z}_+$ and $\mathbb{N}$ be the sets of non-negative integers and positive integers respectively. Denote by f^n the n -th iterate of f ; that is, f^0 is the identity map, and $f^n = f \circ f^{n-1}$ if $n \geq 1$. For any $x \in X$ the subset $O_f(x) = \{f^n(x) : n \in \mathbb{Z}_+\}$ is called the f -orbit of x . A point $x \in X$ is called *periodic* of prime period $n \in \mathbb{N}$ if $f^n(x) = x$ and $f^i(x) \neq x$ for $1 \leq i \leq n-1$. We denote by $Fix(f)$ (resp. $P(f)$) the set of fixed points (resp. periodic points) of f . Let A be a non empty subset of X . It is called *periodic* with period $p \geq 1$ if $A, f(A), \dots, f^{p-1}(A)$ are pairwise disjoint and $f^p(A) = A$. For any $x \in X$, we denote by $\omega_f(x) = \bigcap_{n \geq 0} \overline{O_f(f^n(x))}$ the omega limit set of x . A pair (a, b) in X^2 is called *proximal* if $\liminf_{n \rightarrow +\infty} d(f^n(a), f^n(b)) = 0$, it is called *distal* if $\liminf_{n \rightarrow +\infty} d(f^n(a), f^n(b)) > 0$ and it is called *asymptotic* if $\limsup_{n \rightarrow +\infty} d(f^n(a), f^n(b)) = 0$. A pair (a, b) in X^2 is called a *Li-Yorke* pair (of f) if it is proximal but not asymptotic. A subset S of X with at least two points is a *scrambled* set (of f) if any proper pair $(a, b) \in S^2$ is a Li-Yorke pair. A continuous map $f : X \rightarrow X$ is called *Li-Yorke chaotic* if it has an *uncountable* scrambled set. Denote by $h(f)$ the topological entropy of f (See [1], [7], [10]). For any non empty subset F of a compact metric space X , the set of accumulation points of F , denoted by F' , is called the *derived set* of F . More generally, for any $n \geq 1$, we define $F^{(n)} = (F^{(n-1)})'$ the n -th derivative of F , where $F^0 = F$. If F is closed then F' is a closed

Key words and phrases. Dendrite, dendrite map, ω -limit set, decomposition, Li-Yorke pair, Li-Yorke chaos.

subset of F . A *continuum* is a compact connected metric space. An arc is any space homeomorphic to the compact interval $[0, 1]$. A topological space is *arcwise* connected if any two of its points can be joined by an arc. We use the terminology from Nadler [19]. By a *dendrite* X , we mean a locally connected continuum which contains no homeomorphic copy of a circle i.e simple closed curve. Every sub-continuum of a dendrite is a dendrite ([19], Theorem 10.10) and every connected subset of X is arcwise connected ([19], Proposition 10.9). Let $x \in X$. The number of connected components of $X \setminus \{x\}$, denoted $ord(x, X)$, is called the order of x in X . If $ord(x, X) = 1$ (resp. $ord(x, X) = 2$, resp. $ord(x, X) \geq 3$) then x is called an *endpoint* (resp. *cut point*, resp. *branch point*) of X . If there is no confusion, we denote $ord(x)$ instead of $ord(x, X)$. We denote by $E(X)$ (resp. $B(X)$) the set of endpoints (resp. branch points) of X . A tree, is a dendrite with finitely many endpoints. In addition, any two distinct points x, y of a dendrite X can be joined by a unique arc with endpoints x and y , denote this arc by $[x, y]$ and let $(x, y) = [x, y] \setminus \{y\}$ (resp. $(x, y) = [x, y] \setminus \{x\}$ and $(x, y) = [x, y] \setminus \{x, y\}$). A *free arc* in a dendrite is an arc containing no branch point.

A continuous map from a dendrite into itself is called a *dendrite map*. For any closed subset F of X , we call the *convex hull* of F , noted $[F]$, the intersection of all subdendrites of X containing F .

The ω -limit sets play an important role in studying dynamical systems. Sarkovski [22], proved that if $f : [0, 1] \rightarrow [0, 1]$ is a continuous map with zero topological entropy then any infinite ω -limit set contain no periodic point. This result remain true for graph maps (in particular for tree maps) ([14], Theorem 13). In this paper, we firstly study this question for dendrite maps with closed set of endpoints (See Theorem **A**). Secondly, Smital showed in ([23], Theorem 3.5) that if $L = \omega_f(x)$ is an infinite ω -limit set of an interval map with zero topological entropy (in fact L is uncountable) then there is a sequence $(J_k)_{k \geq 1}$ of f -periodic intervals with the following properties: For any k ,

- (1) J_k has period 2^k ,
- (2) $J_{k+1} \cup f^{2^k}(J_{k+1}) \subset J_k$,
- (3) $L \subset \bigcup_{i=0}^{2^k-1} f^i(J_k)$,
- (4) $L \cap f^i(J_k) \neq \emptyset$ for every $i = 0, 1, \dots, 2^k - 1$.

We will extend this result to dendrite maps $f : X \rightarrow X$ with zero topological entropy, where $E(X)$ is closed and $E(X)'$ finite. This holds in particular if X is a tree. (See Theorem **B**).

In the third part of the paper, we study the question: Does Li-Yorke pair implies chaos for dendrite maps? Kuchta and Smital [17] proved that the existence of Li-Yorke pair implies chaos for interval maps. In [21], Ruette and Snoha proved that the same conclusion holds for graph maps. An example of a triangular map in the square (resp. on the Cantor set and the Warsaw circle) answering negatively the question is found in [11] (resp. [13]). Here

we give some examples of dendrites maps with countable set of endpoints having a Li-Yorke pair but not Li-Yorke chaotic.

Our main results are the following:

Theorem A. Let X be a dendrite with $E(X)$ closed and let $f : X \rightarrow X$ be a dendrite map with zero topological entropy. Let $x \in X$. Then we have

- (1) If $\omega_f(x)$ is infinite then $\omega_f(x) \cap P(f) \subset E(X)'$,
- (2) If $E(X)$ is countable and $\omega_f(x)$ is uncountable then $\omega_f(x) \cap P(f) = \emptyset$.

Remark.

- (1) The condition $\omega_f(x) \cap P(f) \neq \emptyset$ can occur, we built a dendrite map $f : X \rightarrow X$ with zero topological entropy having an infinite ω -limit set $\omega_f(x)$ containing a periodic point, where $E(X)$ is a closed countable set and $\omega_f(x)$ is infinite countable. (See Example 1).
- (2) There is a dendrite map $f : X \rightarrow X$ with zero topological entropy having an ω -limit set $\omega_f(x)$ containing a periodic point with $E(X)$ non closed and countable and $\omega_f(x)$ uncountable. (See Example 2).
- (3) There is a dendrite map $f : X \rightarrow X$ with zero topological entropy having an uncountable ω -limit set $\omega_f(x)$ containing a periodic point with $E(X)$ closed and uncountable. (See Example 3).

Theorem B. Let X be a dendrite such that $E(X)$ is closed set having finitely many accumulation points and let $f : X \rightarrow X$ be a dendrite map with zero topological entropy. Let L be an uncountable ω -limit set. Then there is a sequence of f -periodic subdendrites $(D_k)_{k \geq 1}$ of X and a sequence of integers $n_k \geq 2$ for every $k \geq 1$ with the followings properties: $\forall k \geq 1$,

- (1) D_k has period $\alpha_k := n_1 n_2 \dots n_k$,
- (2) $\cup_{j=0}^{n_k-1} f^{j\alpha_k}(D_k) \subset D_{k-1}$; $j \geq 2$,
- (3) $L \subset \cup_{i=0}^{\alpha_k-1} f^i(D_k)$,
- (4) $f(L \cap f^i(D_k)) = L \cap f^{i+1}(D_k)$; $0 \leq i \leq \alpha_k - 1$. In particular $L \cap f^i(D_k) \neq \emptyset$, $\forall 0 \leq i \leq \alpha_k - 1$,
- (5) $\forall 0 \leq i \neq j < \alpha_k$, $f^i(D_k) \cap f^j(D_k)$ has empty interior.

Corollary 1. Let X be a tree and $f : X \rightarrow X$ a continuous map with zero topological entropy. Let $L = \omega_f(x)$ an infinite ω -limit set. Then there is a sequence $(J_k)_{k \geq 1}$ of f -periodic arcs and a sequence of integers $n_k \geq 2$ for every $k \geq 1$ with the following properties: For any k ,

- (1) J_k has period $\alpha_k := n_1 n_2 \dots n_k$,
- (2) $\cup_{j=0}^{n_k-1} f^{j\alpha_k}(J_k) \subset J_{k-1}$; $j \geq 2$,
- (3) $L \subset \cup_{i=0}^{\alpha_k-1} f^i(J_k)$,
- (4) $f(L \cap f^i(J_k)) = L \cap f^{i+1}(J_k)$; $0 \leq i \leq \alpha_k - 1$. In particular $L \cap f^i(J_k) \neq \emptyset$, $\forall 0 \leq i \leq \alpha_k - 1$,
- (5) $\forall 0 \leq i \neq j < \alpha_k$, $f^i(J_k) \cap f^j(J_k)$ are either disjoint or they intersect in their common endpoints.

Theorem C. Let X be a dendrite with $E(X)$ closed and $E(X)'$ finite. Let $f : X \rightarrow X$ be a dendrite map. If f has a Li-Yorke pair (x, y) such that

$\omega_f(x)$ or $\omega_f(y)$ is uncountable then f is Li-Yorke chaotic.

Remark. Theorem **C** is not always true if X is a dendrite with $E(X)$ closed countable with $E(X)'$ infinite. (See Example 4).

2. Preliminaries

Lemma 2.1. ([18], Lemma 2.1) *Let (X, d) be a dendrite. Then, for every $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that, for any $x, y \in X$ with $d(x, y) \leq \delta$, the diameter $\text{diam}([x, y]) < \varepsilon$.*

Here $\text{diam}(A) := \sup_{x, y \in A} d(x, y)$ where A is a non empty subset of (X, d) .

Corollary 2.2. *Let (X, d) be a dendrite. Then, for every $x, y \in X; x \neq y$, there is $\varepsilon > 0$ and an arc $J \subset [x, y]$ such that for any $u \in B(x, \varepsilon)$ and $v \in B(y, \varepsilon)$ we have $J \subset [u, v]$.*

Proof. Fix $x, y \in X; x \neq y$. Let U and V two disjoint subdendrites of X such that $x \in \text{int}(U)$ and $y \in \text{int}(V)$ (where $\text{int}(A)$ denote the interior of the subset A). Since U, V and $[x, y]$ are connected then by Theorem 10.10 of [19], $U \cap [x, y]$ and $V \cap [x, y]$ are also connected. Let $c_1, c_2 \in X$ satisfying $U \cap [x, y] = [x, c_1]$ and $V \cap [x, y] = [c_2, y]$. We have $[x, c_1] \cap [y, c_2] \subset U \cap V = \emptyset$ so $[x, c_1] \cap [y, c_2] = \emptyset$.

Let $u \in U$ and $v \in V$ arbitrarily and denote by $u' = r_{[x, y]}(u), v' = r_{[x, y]}(v)$ where $r_{[x, y]}$ is the first point map for $[x, y]$ (See [19], page 176). We have $u' \in [x, u] \cap [x, y] \subset U \cap [x, y] = [x, c_1]$. Similarly, $v' \in [y, c_2]$. Since $[u, u'] \cap [x, y] = [v, v'] \cap [x, y] = \emptyset$ (If $a = b$ then $[a, b] = \emptyset$) then $[u, u'], [v, v']$ and $[u', v']$ are pairwise disjoint so $[u, v] = [u, u'] \cup [u', v'] \cup [v', v]$. Finely, since $u' \in [x, c_1], v' \in [c_2, y]$ then $[u', v']$ contains $J := [c_1, c_2]$ (which is independent from u, v) so $J \subset [u, v]$. By taking $\varepsilon > 0$ with $B(x, \varepsilon) \subset U$ and $B(y, \varepsilon) \subset V$, we finish the proof of the corollary. $\square$

Lemma 2.3. *Let X be a dendrite and C_1, C_2 two disjoint connected subsets in X . Then $\overline{C_1} \cap \overline{C_2}$ is at most one point.*

Proof. Suppose that $\overline{C_1} \cap \overline{C_2}$ contains at least two distinct points a, b . Then by Theorem 10.10 of [19] this intersection is connected, so it is arcwise connected. Hence we obtain $[a, b] \subset \overline{C_1} \cap \overline{C_2}$. There is four sequences $(a_n)_{n \geq 1}, (b_n)_{n \geq 1}$ in C_1 and $(a'_n)_{n \geq 1}, (b'_n)_{n \geq 1}$ in C_2 such that $(a_n)_{n \geq 1}, (a'_n)_{n \geq 1}$ converges to a and $(b_n)_{n \geq 1}, (b'_n)_{n \geq 1}$ converges to b . By Corollary 2.2, there is $\varepsilon > 0$ and an arc $J \subset [a, b]$ such that $J \subset [u, v]$ for any $u \in B(a, \varepsilon)$ and $v \in B(b, \varepsilon)$. There is $n > 0$ such that $a_n, a'_n \in B(a, \varepsilon)$ and $b_n, b'_n \in B(b, \varepsilon)$. So we have $J \subset [a_n, b_n] \cap [a'_n, b'_n]$. Since C_1 and C_2 are arcwise connected then $[a_n, b_n] \subset C_1$ and $[a'_n, b'_n] \subset C_2$ then $J \subset C_1 \cap C_2$, absurd. $\square$

Lemma 2.4. *Let X be a dendrite and F a non empty closed subset of X . Let $a \in F$, then*

- (1) $[F] = \cup_{z \in F} [a, z]$,
 (2) $E([F]) \subset F$ and we have $E([F]) = F$ when $F \subset E(X)$.

Proof. (1) For any $z \in F$, $[a, z]$ is connected then the subset $G := \cup_{z \in F} [a, z]$ is connected and contains F . We will prove that G is closed. Let $(w_n)_{n \geq 1}$ be a sequence in G converging to $w \in X$. For any $n \geq 1$, there is $z_n \in F$ such that $w_n \in [a, z_n]$. Since F is compact, by considering a sub sequence of $(z_n)_{n \geq 1}$, we may assume that $(z_n)_{n \geq 1}$ converges to $z \in F$. It suffices to prove that $w \in [a, z]$. Suppose contrarily that $\delta := d(w, [a, z]) > 0$ then $B(w, \frac{\delta}{2}) \cap B([a, z], \frac{\delta}{2}) = \emptyset$. By Lemma 2.1, since $(z_n)_{n \geq 1}$ converges to z then there is $N > 0$ such that $\text{diam}([z, z_n]) < \frac{\delta}{2}$ when $n > N$, hence $[z, z_n] \subset B(z, \frac{\delta}{2})$ so $[a, z_n] \subset [a, z] \cup [z, z_n] \subset B([a, z], \frac{\delta}{2})$. Let $n > N$ such that $d(w, w_n) < \frac{\delta}{2}$. Then $w_n \in B([a, z], \frac{\delta}{2}) \cap B(w, \frac{\delta}{2})$ hence $B([a, z], \frac{\delta}{2}) \cap B(w, \frac{\delta}{2}) \neq \emptyset$, absurd. So $w \in G$ then we conclude that G is a subdendrite of X satisfying $F \subset G \subset [F]$, hence $[F] = G$.

(2) Let $e \in E([F])$, by (1) there is $z \in F$ such that $e \in [a, z]$. Since $1 \leq \text{ord}(e, [a, z]) \leq \text{ord}(e, [F]) = 1$ hence $e = z$ or a then $e \in F$, so $E([F]) \subset F$. Now, suppose that $F \subset E(X)$. It suffices to prove that $F \subset E([F])$. Since $F \subset [F]$ then $F \subset [F] \cap E(X) \subset E([F])$. This finish the proof of this Lemma. □

Lemma 2.5. ([18], Lemma 2.3) *Let $(C_n)_{n \geq 1}$ be a sequence of pairwise disjoint connected subsets of a dendrite (X, d) . Then we have $\lim_{n \rightarrow +\infty} \text{diam}(C_n) = 0$.*

Lemma 2.6. ([3], Theorem 3.3) *If a dendrite has a closed set of endpoints then any point of it has finite order.*

Theorem 2.7. ([3], Theorem 3.2) *Every subcontinuum of a dendrite with a closed set of endpoints is a dendrite with a closed set of endpoints.*

Proposition 2.8. ([3], Corollary 3.5) *If X is a dendrite with $E(X)$ closed set then $\overline{B(X)} \subset B(X) \cup E(X)$. In particular, $B(X) \cup E(X)$ is closed.*

Proposition 2.9. ([3], Corollary 3.6) *If X is a dendrite with $E(X)$ closed then $B(X)$ is a discrete set.*

Lemma 2.10. ([8], Proposition 4.14) *If Y and X dendrites with $Y \subset X$, then $E(Y)' \subset E(X)'$.*

Lemma 2.11. ([20], Lemma 4) *If Y and X are dendrites with $Y \subset X$, then $\text{card}(E(Y)) \leq \text{card}(E(X))$.*

For a subset A of X , we denote by $\text{card}(A)$ the number of elements of A . If $A = \emptyset$, we take $\text{card}(A) = 0$.

Definition 2.12. *Let $f : X \rightarrow X$ be a dendrite map and I, J two arcs in X . We say that I, J form an arc horseshoe for f if $f^n(I) \cap f^m(J) \supset I \cup J$ for some $n, m \in \mathbb{N}$, where I, J have exactly a common one endpoint.*

Theorem 2.13. [5] *Let X be a compact metric space and $f : X \rightarrow X$ a continuous map. If $h(f) > 0$ then f is Li-Yorke chaotic.*

When X is a compact interval, it is well known that if f^n has an arc horseshoe for some $n \in \mathbb{N}$, then $h(f) > 0$. Actually, for dendrite map, we have

Theorem 2.14. ([16], Theorem 2) *Let $f : X \rightarrow X$ be a dendrite map. If f has an arc horseshoe then $h(f) > 0$.*

Lemma 2.15. ([6], page 71) *Let (X, d) be a compact metric space and $f : X \rightarrow X$ a continuous map. Let F be a proper closed subset of an ω -limit set $L = \omega_f(x)$, then*

$$\overline{f(L \setminus F)} \cap F \neq \emptyset.$$

Lemma 2.15 is equivalent to the following Lemma:

Lemma 2.16. *If G is a non empty open subset of L (relatively to L) and such that $f(\overline{G}) \subset G$ then $G = L$.*

Proof. Suppose contrarily that there is a non empty open subset G of L such that $f(\overline{G}) \subset G$ and $G \neq L$. Then $F := L \setminus G$ is a non empty closed subset of L and $\overline{f(L \setminus F)} \cap F = \overline{f(G)} \cap F = f(\overline{G}) \cap F \subset G \cap F = \emptyset$, absurd. $\square$

3. On ω -limit set containing a periodic point.

To prove Theorem A, we need the following Lemmas

Lemma 3.1. ([9], Proposition 4.4) *Let X be a dendrite with $E(X)$ closed and $E(X)' \neq E(X)$. Then for any $e \in E(X) \setminus E(X)'$, there is $b \in X, b \neq e$ such that $[e, b]$ is a neighborhood of e and $[e, b] \cap B(X) = \emptyset$.*

Lemma 3.2. *Let X be a dendrite with countable closed set of endpoints and let Y be a subdendrite of X . Denote by $Y_1 := Y \setminus [B(X) \cup E(Y)] = \cup_{i \in I} J_i$, where $(J_i)_{i \in I}$ is the sequence of the connected components of Y_1 . Then I is at most countable and each J_i is an open free arc in X .*

Proof. Set $F = (B(X) \cap Y) \cup E(Y)$. We will show that $Y_1 = Y \setminus F$ is closed in Y : Indeed, by theorem 2.7, since $E(X)$ is closed, so $E(Y)$ is also closed. Let $(b_n)_{n \geq 1}$ be an infinite sequence in $B(X) \cap Y$ converging to b . Then $b \in Y$ and by Proposition 2.8, $b \in B(X) \cup E(X)$, hence $b \in (Y \cap B(X)) \cup (Y \cap E(X)) \subset (Y \cap B(X)) \cup E(Y)$ so F is closed in Y , hence Y_1 is open in Y . So by ([19], p. 120), each component J_i is open in Y . For any $i \in I$, J_i is open in X , since $J_i \cap B(X) = \emptyset$.

Let us prove that I is at most countable. For any $i \in I$, write $\overline{J_i} = [a_i, b_i]$ and define the map $h : I \rightarrow (B(X) \cup E(Y))^2$ as follow: $\forall i \in I, h(i) = \{a_i, b_i\}$. The map h is well defined (since X is uniquely arcwise connected) and it is one-to-one then I is at most countable since $B(X) \cup E(Y)$ is at most countable. $\square$

Lemma 3.3. *Let $f : X \rightarrow X$ be a dendrite map such that $E(X)$ is closed and countable. Let $a \in \text{Fix}(f)$ and $L := \omega_f(x)$ an uncountable ω -limit set such that $L \cap P(f) = \emptyset$ then for any $y \in L$, there is $p, k \geq 0$ such that $[a, f^k(x)] \subset [a, f^p(y)]$.*

Proof. Let $y \in L$. We have $\omega_f(y)$ is a closed invariant subset by f then there is a minimal subset, denoted by K , in $\omega_f(y)$. Since $L \cap P(f) = \emptyset$ then K has no periodic point. So K is infinite and has no isolated point, hence K is uncountable so it is for $\omega_f(y)$. Now, denote by $(C_i)_{i \in \mathbb{N}}$ the sequence of connected components of $X \setminus (B(X) \cup E(X))$. By Lemma 3.2, each C_i is an open free arc in X . There is $j \in \mathbb{N}$ such that $\omega_f(y) \cap C_j$ is uncountable. Let $u, v \in \omega_f(y) \cap C_j$ such that $u \in (a, v)$. There is two open disjoint arcs I_u, I_v in C_j such that $u \in I_u, v \in I_v$. Let $p, k' \geq 0$ such that $f^p(y) \in I_v$ and $f^{k'}(y) \in I_u$, since $f^{k'}(y) \in L$, there is $k > 0$ such that $f^k(x) \in I_u$, so we obtain the inclusion $[a, f^k(x)] \subset [a, f^p(y)]$. $\square$

Proof of Theorem A.

(1) Denote by $L = \omega_f(x)$. Suppose that $L \cap P(f) \not\subset E(X)'$, there is $a \in L \setminus E(X)'$ such that $f^N(a) = a$ for some $N > 0$. Since L is infinite and $\forall 0 \leq i \leq N-1, f(\omega_{f^N}(f^i(x))) = \omega_{f^N}(f^{i+1}(x))$ then $\omega_{f^N}(f^i(x))$ is infinite for any $0 \leq i \leq N-1$. Let $0 \leq j \leq N-1$ such that $a \in \omega_{f^N}(f^j(x))$, so we may assume that $a \in \text{Fix}(f)$. By Corollary 2.6, since $E(X)$ is closed then $1 \leq n := \text{ord}(a) < +\infty$. By Corollary 2.9 and Lemma 3.1, a has a neighborhood, V , which is a tree such that $V \cap B(X) \subset \{a\}$. We can write $V = \cup_{i=1}^n [a, b_i]$ such that the subsets $(a, b_i]; 1 \leq i \leq n$ are pairwise disjoint.

Claim 1. *There is $1 \leq i_0 \leq n$ and an infinite sequence of periodic points in (a, b_{i_0}) converging to a .*

By ([6], Lemma 4) a is not isolated relatively to L , then there is an infinite sequence in L , say $(y_n)_{n \geq 1}$, converging to a . Since V is a neighborhood of a then we may assume that $(y_n)_{n \geq 1} \subset V$. Let $1 \leq i_0 \leq n$ such that $a \in \overline{(a, b_{i_0})} \cap \overline{(y_n)_{n \geq 1}}$. By considering a sub sequence we may assume that $(y_n)_{n \geq 1} \subset (a, b_{i_0}]$. Let $c \in (a, b_{i_0})$ arbitrarily. There is $1 \leq n_1 < n_2 < n_3$ such that $y_{n_2} \in (y_{n_1}, y_{n_3}) \subset (a, c)$. Let I_1, I_2 and I_3 a disjoint open arc in (a, c) such that $y_{n_i} \in I_i; i = 1, 2, 3$. There is $n, m \geq 0$ and $p, q > 0$ such that $f^n(x) \in I_1, f^m(x) \in I_3$ and $f^p(f^n(x)), f^q(f^m(x)) \in I_2$. So we have $\{f^p(f^n(x)), f^q(f^m(x))\} \subset (f^n(x), f^m(x))$, then by [2] (one can use also Theorem 2.13 of [18]), $P(f) \cap (f^n(x), f^m(x)) \neq \emptyset$ hence $P(f) \cap (a, c) \neq \emptyset$. This finish the proof of Claim 1.

Now, denote by $(C_k)_{1 \leq k \leq n}$ the sequence of connected components of $X \setminus \{a\}$ such that $b_k \in C_k, \forall 1 \leq k \leq n$ and let $c \in \text{Fix}(f^r) \cap (a, b_{i_0}); r \geq 1$ such that $L \cap (C_{i_0} \setminus [a, c]) \neq \emptyset$. Denote by $g = f^r$. There is $n \geq 0$ such that $f^n(x) \in (a, c)$, since $L = \omega_f(f^n(x))$, we may assume that $n = 0$. We distinguish two cases both of them lead us to a contradiction:

Case 1. If $O_g(x) \not\subseteq C_{i_0}$. Let $1 \leq j \leq n; j \neq i_0$ and $p, k > 0$ such that $g^p(x) \in C_j$ and $f^k(x) \in C_{i_0} \setminus [a, c]$. Let $I = [a, x]$ and $J = [x, c]$. Then we have $f^k(I) \supset [a, f^k(x)] \supset [a, c] = I \cup J$ and $f^{rp}(J) = g^p(J) \supset [c, g^p(x)] \supset [a, c] = I \cup J$, so I, J form an arc horseshoe then by [16] and [4] we have $h(f) > 0$, absurd.

Case 2. If $O_g(x) \subseteq C_{i_0}$. Denote by $F_c = \cup_{n=0}^{+\infty} g^{-n}(c) \cap [a, c]$.

(a) If there is $z \in F_c \setminus \{c\}$ such that $(c, z] \cap O_g(x) \neq \emptyset$. Let $n \geq 0, k > 0$ such that $g^n(x) \in (c, z)$ and $g^k(z) = c$. Let $I = [c, g^n(x)]$ and $J = [z, g^n(x)]$. Since $a \in \omega_g(x)$ then there is $p > k$ such that $I \cup J \subset [c, g^{n+p}(x)]$. So we have $g^p(I) \supset [c, g^{n+p}(x)] \supset I \cup J$ and $g^k(J) \supset [c, g^{n+k}(x)] \supset I \cup J$, hence $g^p(J) = g^{p-k}(g^k(J)) \supset [c, g^{n+p}(x)] \supset I \cup J$ so I, J form an arc horseshoe for g , hence $h(f) = \frac{1}{r}h(g) > 0$, absurd.

(b) If for any $z \in F_c; [c, z] \cap O_g(x) = \emptyset$.

Claim 2. We have $O_g(x) \subset [a, c]$.

Since otherwise, there is $n > 0$ such that $g^n(x) \notin [a, c]$. (i) If $g^n(x) \in C_{i_0} \setminus [a, c]$ then $g^n([a, x]) \supset [a, g^n(x)] \ni c$, so there is $c_{-1} \in (a, x); g^n(c_{-1}) = c$, hence $x \in [c, c_{-1}] \cap O_g(x)$, a contradiction. (ii) If $g^n(x) \in C_j$ for some $j \neq i_0$. Then $g^n([c, x]) \supset [c, g^n(x)] \ni a$. Let $a_{-1} \in (c, x); g^n(a_{-1}) = a$ and $k > 0$ such that $f^k(x) \in C_{i_0} \setminus [a, c]$. Then $f^{rn}([c, a_{-1}]) = g^n([c, a_{-1}]) \supset [a, c]$ and $f^k([a, a_{-1}]) \supset f^k([a, x]) \supset [a, f^k(x)] \supset [a, c]$, so $[a, a_{-1}]$ and $[c, a_{-1}]$ form an arc horseshoe for f , absurd. This finish the proof of Claim 2.

Now, we have $\omega_g(x) \subset [a, c]$. Denote by $[a, d] = [\omega_g(x)]$ the convex hull of $\omega_g(x)$. We remark that $d \in \omega_g(x)$ since $\omega_g(x)$ is closed.

Claim 3. For any $n \geq 0, g^n([a, d]) \subset [a, c]$.

Suppose contrarily that there is $n > 0$ such that $g^n([a, d]) \not\subseteq [a, c]$. Let $z \in (a, d), g^n(z) \notin [a, c]$. (i) If $g^n(z) \in C_{i_0} \setminus [a, c]$ then c has an antecedent c_{-1} by g^n in $(a, z]$, since $d \in [c, c_{-1}] \cap \omega_g(x)$ then $O_g(x) \cap (c, c_{-1}) \neq \emptyset$, a contradiction. (ii) If $g^n(z) \in C_j$ for some $j \neq i_0$. Then $g^n([c, z]) \supset [c, g^n(z)] \ni a$ then there is $a_{-1} \in (c, z]$ such that $g^n(a_{-1}) = a$. Let $k, p > 0$ such that $f^k(x) \in (a, a_{-1}), f^{k+p}(x) \in C_{i_0} \setminus [a, c]$. We have $f^p([a, f^k(x)]) \supset [a, f^{k+p}(x)] \supset [a, f^k(x)] \cup [f^k(x), c]$ and $f^{nr}([c, f^k(x)]) \supset f^{nr}([c, a_{-1}]) = g^n([c, a_{-1}]) \supset [c, a] = [c, f^k(x)] \cup [f^k(x), a]$, hence $[c, f^k(x)], [a, f^k(x)]$ form an arc horseshoe for f , absurd. This finish the proof of the Claim 3.

Now, we have $g^n([a, d]) = g^n([a, d]) \cup g^n(d) \subset [a, c] \cup \omega_g(x) \subset [a, c]$. In the other hand, since $[a, d] \supset \omega_g(x)$ so $g([a, d]) \supset [a, d]$. The subset $I = \cup_{n=0}^{+\infty} g^n([a, d])$ is connected included in $[a, c]$ and strongly invariant by g so it is for $J = \bar{I} \subset [a, c]$. Now $g|_J : J \rightarrow J$ is a continuous interval map with $\omega_{g|_J}(x) = \omega_g(x); x \in J$ is infinite containing a fixed point a , so

by [22], $h(g) \geq h(g|_J) > 0$ hence $h(f) > 0$, absurd. We conclude that $L \cap P(f) \subset E(X)'$.

(2) Now, suppose that $L = \omega_f(x)$ uncountable and $E(X)$ countable. By (1) of this Theorem, we have $L \cap P(f) \subset E(X)'$. Suppose that there is $a \in L \cap P(f)$, we may assume that $a \in \text{Fix}(f)$. By Lemma 3.2, write $X \setminus [B(X) \cup E(X)] = \cup_{i=1}^{+\infty} J_i$ where each J_i is an open free arc in X . There is $i_0 > 0$ such that $L \cap J_{i_0}$ is uncountable. Write $J_{i_0} = (u, v)$ such that $v \in (a, u)$. There exists $c \in \text{Fix}(f^r)$; $r \geq 1$ such that $L \cap (u, c)$ is uncountable. Denote by $g = f^r$. There is $0 \leq i \leq r-1$ such that $\omega_g(x_i) \cap (u, c)$ is uncountable; $x_i = f^i(x)$. There is $k \geq 0, p > 0$ such that $g^k(x_i), g^{k+p}(x_i) \in (w, c) \cap (u, c)$ for some $w \in \omega_g(x_i)$. Denote by X_0, X_1 the connected components of $X \setminus \{c\}$ such that $a \in X_1$ and let $l_j = X_j \cap \omega_g(x_i)$; $j = 0, 1$. By (1), $c \notin \omega_g(x_i)$ then l_0 and l_1 are two non empty clopen sets relatively to $\omega_g(x_i)$, hence by Lemma 2.16 we have $\forall n \geq 1, g^n(l_0) \not\subset l_0$. Let $y \in l_0$ such that $g^p(y) \in l_1$.

We will build an arc horseshoe. We distinguish two cases:

Case 1. *There is an infinite sequence $(n_l)_{l>0}$ such that $(g^{n_l}(x_i))_{l>0}$ converges to y and $g^{n_l}(x_i) \in (c, y)$; $\forall l > 0$.* By continuity of g^p , there is $n > 0$ such that $g^n(x_i) \in X_0, g^{n+p}(x_i) \in X_1$. Denote by $I = [c, g^k(x_i)], J = [g^k(x_i), g^n(x_i)]$ if $g^k(x_i) \in (c, g^n(x_i))$, (resp. $I = [c, g^n(x_i)], J = [g^n(x_i), g^k(x_i)]$ if $g^n(x_i) \in (c, g^k(x_i))$). We have $g^p(I) \cap g^p(J) \supset [c, g^{k+p}(x_i)]$ (respectively, $g^p(I) \cap g^p(J) \supset [c, g^{n+p}(x_i)]$). There is $s > 0$ such that $[c, g^{k+p+s}(x_i)] \supset I \cup J$ hence $g^{p+s}(I) \cap g^{p+s}(J) \supset I \cup J$ (resp. since $g^k(x_i) \in (c, w)$; $w \in \omega_g(x_i)$ then there is $r > 0$ such that $[c, g^{n+p+r}(x_i)] \supset I \cup J$ hence $g^{p+r}(I) \cap g^{p+r}(J) \supset I \cup J$). So I, J form an arc horseshoe for f , absurd.

Case 2. *There is an infinite sequence $(n_l)_{l>0}$ such that $(g^{n_l}(x_i))_{l>0}$ converges to y and $y \in (c, g^{n_l}(x_i))$; $\forall l > 0$.* Similarly as in Case 1, we build an arc horseshoe by theses three points $c, g^k(x_i), y$ (resp. by $c, g^n(x_i), g^k(x_i)$ for a convenient integer n) if $g^k(x_i) \in (c, y)$ (resp. if $y \in (c, g^k(x_i))$).

This finish the proof of Theorem A. $\square$

4. Examples of dendrite maps with zero topological entropy.

4.1. Example 1. We build a dendrite X with $E(X)$ countable closed set and a map $f : X \rightarrow X$ with zero topological entropy having an infinite ω -limit set containing a periodic point.

Construction of the dendrite X . For any $n \geq 0$, let

- (1) $a_n = (1 - \frac{1}{n+1}, 0), b_n = (1 - \frac{1}{n+1}, \frac{(-1)^n}{n+1})$ and $e = (1, 0)$,
- (2) $w_n \in (a_n, b_n)$,
- (3) $(b_n^k)_{k \geq 0}$ be a monotone sequence in $[b_n, w_n)$ converging to w_n where $b_n^0 = b_n$,
- (4) $X = \cup_{i=0}^{+\infty} [a_i, b_i] \cup [a_0, e]$.

We can see that X is a dendrite with $E(X) = \{b_n; n \geq 0\} \cup \{e\}$ closed and $E(X)' = \{e\}$.

Construction of the map $f : X \rightarrow X$. (See Figure 1) We define f as follows: For any $n, k \in \mathbb{Z}_+$,

- (1) $f(e) = e$,
 f maps linearly
- (2) $[a_n, a_{n+1}]$ to $[a_{n+1}, a_{n+2}]$ such that $f(a_n) = a_{n+1}$,
- (3) $[a_{2n+2}, w_{2n+2}]$ to $[a_{2n+3}, w_{2n}]$,
- (4) $[a_{2n+1}, w_{2n+1}]$ to $[a_{2n+2}, w_{2n+3}]$,
- (5) $[a_0, w_0]$ to $[a_1, w_1]$,
- (6) $[w_{2n+2}, b_{2n+2}]$ to $[w_{2n}, b_{2n}]$ such that $f(b_{2n+2}^k) = b_{2n}^{k+1}$,
- (7) $[w_0, b_0]$ to $[w_1, b_1]$ such that $f(b_0^k) = b_1^k$,
- (8) $[w_{2n+1}, b_{2n+1}^1]$ to $[w_{2n+3}, b_{2n+3}]$ such that $f(b_{2n+1}^{k+1}) = b_{2n+3}^k$,
- (9) $[b_{2n+1}, b_{2n+1}^1]$ to $[b_{2n+3}, b_{2n+2}]$.




FIGURE 1. Dendrite with $E(X)'$ is reduced to one point.

Denote by $x = b_0$ and for any $n \geq 0, x_n = f^n(x)$. Then the first fifteen elements of the orbit of x are:

$$x = b_0, x_1 = b_1, x_2 = b_2, x_3 = b_0^1, x_4 = b_1^1, x_5 = b_3, x_6 = b_4, x_7 = b_2^1, x_8 = b_0^2, x_9 = b_1^2, x_{10} = b_3^1, x_{11} = b_5, x_{12} = b_6, x_{13} = b_4^1, x_{14} = b_2^2, x_{15} = b_0^3.$$

Lemma 4.1. *The map $f : X \rightarrow X$ obtained is continuous and satisfy the following properties:*

- (1) $\omega_f(b_0) = \{w_n; n \geq 0\} \cup \{e\}$. Therefore (b_0, e) is a Li-Yorke pair,

- (2) for any $y \in X$, we have either $\omega_f(y) = \{e\}$ or $\omega_f(y) = \omega_f(b_0)$,
- (3) f has zero topological entropy,
- (4) (X, f) is proximal i.e any pair $(x, y) \in X^2$ is proximal.

Proof. It is easy to prove (1) and (2). Let prove (3). Denote by $R(f)$ the set of recurrent points of f i.e $R(f) := \{x \in X, x \in \omega_f(x)\}$. Since $E(X)$ is countable then by [18] we have $\overline{R(f)} = \overline{P(f)}$. Since $P(f) = \{e\}$ then $\overline{R(f)} = \{e\}$. Now by [6], page 196 we have $h(f) = h(f|_{\overline{R(f)}}) = h(f|_{\{e\}}) = 0$.
 (4) Let prove that (X, f) is proximal. Let $(x, y) \in X^2$. Fix $\varepsilon > 0$. There is $L > 0$ such that for any $i \geq 0$, $\{f^i(y), f^{i+1}(y), \dots, f^{i+L}(y)\} \cap B(e, \frac{\varepsilon}{2}) \neq \emptyset$. Since $e = f(e) \in \omega_f(x)$ there is $p > 0$ such that $\{f^p(x), f^{p+1}(x), \dots, f^{p+L}(x)\} \subset B(e, \frac{\varepsilon}{2})$. Let $0 \leq k \leq L$ such that $f^{p+k}(y) \in B(e, \frac{\varepsilon}{2})$, we obtain $d(f^{p+k}(x), f^{p+k}(y)) < \varepsilon$. □

4.2. Example 2. We will prove that there is a non chaotic dendrite map $f : X \rightarrow X$ with $E(X)$ countable but non closed set having an uncountable ω -limit set containing a periodic point.

Construction of the dendrite X .

For any $n \geq 1$, denote by:

- $A = (0, 0)$ and $B = (1, 0)$,
- $S_n = \{\frac{i}{2^n} \text{ where } 1 \leq i \leq 2^n \text{ is odd}\}$
- $S_{2n+1} = \{a_{2^{2n}} > a_{2^{2n+1}} > \dots > a_{2^{2n+1-1}}\}; n \geq 0$ and $S_{2n} = \{a_{2^{2n-1}} < a_{2^{2n-1+1}} < \dots < a_{2^{2n-1}}\}$,
- $I_k = [A_k, B_k]$ where $A_k = (a_k, 0)$ and $B_k = (a_k, \frac{1}{n+1})$ for any $n \geq 0$ and $k \in \{2^n, 2^n + 1, \dots, 2^{n+1} - 1\}$.

So the set $X := [A, B] \cup (\cup_{k \geq 1} I_k)$ is a dendrite with $E(X) = \{B_k; k \geq 1\}$ and $\overline{E(X)} = E(X) \cup [A, B]$.

Construction of the map $f : X \rightarrow X$. (See Figure 2).

The map f is defined as follow: f fix any point in $[A, B]$ and for any $k \geq 1$, f maps linearly I_k to I_{k+1} such that the center of I_k is sent to A_{k+1} .

The map $f : X \rightarrow X$ satisfies the following properties: for any $k \geq 0$, $\omega_f(B_k) = [A, B]$ and for any $y \in X \setminus E(X)$, there is $p \geq 0$ such that $f^p(y) \in [A, B]$. It follows that $(u, v) \in X^2$ is a Li-Yorke pair if and only if either (u, v) or (v, u) lies to $E(X) \times X \setminus E(X)$. Then f is not chaotic so $h(f) = 0$ but $\omega_f(B_0) = [A, B] = \text{Fix}(f)$.




FIGURE 2. Dendrite with a non closed countable set of endpoints.

4.3. Example 3.

Lemma 4.2. [12] *There is $s \in \Sigma_2 := \{0, 1\}^{\mathbb{Z}_+}$ a recurrent point such that $\omega_\sigma(s)$ is an uncountable ω -limit set containing a fixed point and $\sigma|_{\omega_\sigma(s)}$ has zero topological entropy, where $\sigma : \Sigma_2 \rightarrow \Sigma_2$ is the shift map defined as follow: $\forall x = (x_n)_{n \geq 0} \in \Sigma_2, \sigma(x) = (x_{n+1})_{n \geq 0}$.*

Lemma 4.3. ([3], Proposition 6.8, p. 16) *Each dendrite with an uncountable set of its endpoints contains a homeomorphic copy of the Gehman dendrite.*

Proposition 4.4. *Let X be a dendrite with $E(X)$ uncountable. Then there is a continuous map $f : X \rightarrow X$ with zero topological entropy having an uncountable ω -limit set containing a periodic point.*

Proof. Let X be a dendrite with $E(X)$ uncountable. If X is a Gehman dendrite. The set $E := E(X)$ is homeomorphic to $\omega_\sigma(s)$ where s is defined in the Lemma 4.2. Let f be the map defined in [15] with the same notations such that f act in E as the subshift $\sigma_1 := \sigma|_{\omega_\sigma(s)}$. We may assume that $E = \omega_\sigma(s)$. Any point $y \in X \setminus E$ is eventually mapped to c so $\overline{R(f)} \subset E \cup \{c\}$. Since $h(f) = h(f|_{\overline{R(f)}})$ then $h(f) \leq \max(h(f|_E), h(f|_c)) = h(f|_E) = h(\sigma_1) = 0$ hence $h(f) = 0$. Also we have $\omega_f(s) = \omega_\sigma(s)$ is uncountable containing a periodic point. Generally, by Lemma 4.3, X contains a homeomorphic copy of the Gehman dendrite, G . Denote by $r_G : X \rightarrow G$ the retraction map. Let $f : G \rightarrow G$ defined above and we set $g = f \circ r_G : X \rightarrow X$. Then we have g is a dendrite map on X , $h(g) = h(f)$ and there is $x \in G$ such that $\omega_g(x) = \omega_f(x)$ is uncountable containing a periodic point. This finish the proof of the Proposition. $\square$

5. Decomposition of an uncountable ω -limit set.

Assumption 5.1. *We assume that X is a dendrite with $E(X)$ closed and $E(X)'$ finite. Let $f : X \rightarrow X$ be a dendrite map with zero topological entropy having an uncountable omega limit set $L := \omega_f(x)$ (such map exists by ([23], Theorem 2.7)). Denote by $M = [L]$ the convex hull of L .*

The aim of this paragraph is to prove the following proposition.

Proposition 5.2. *Let Assumption 5.1 be satisfied. Then there is an integer $n \geq 2$ and a connected subset S of X such that:*

- (i) $S, f(S), \dots, f^{n-1}(S)$ are pairwise disjoint,
- (ii) $f^n(S) = S$,
- (iii) $L \subset \bigcup_{i=0}^{n-1} f^i(S)$,
- (iv) $\forall 0 \leq i \leq n-1, f(L \cap f^i(S)) = L \cap f^{i+1}(S)$.

Remark 5.3. *In fact, (iv) is an immediate consequence of (i), (ii) and (iii).*

Remark 5.4. *If $f : X \rightarrow X$ is a tree map with zero topological entropy, then any infinite ω -limit set is uncountable.*

Lemma 5.5. *Let assumption 5.1 be satisfied. Let Y be a (non degenerate) subdendrite of X such that $E(Y) \cap E(X)' = \emptyset$, then Y is a tree and $X \setminus Y$ has finitely many connected components. Furthermore, there is a pairwise disjoint subdendrites $D_1, D_2, \dots, D_n$ in X such that $X \setminus Y \subset \bigcup_{i=1}^n D_i$; $n > 0$ and for any $1 \leq i \leq n$, $D_i \cap Y$ is reduced to one point.*

Proof. Suppose contrarily that $X \setminus Y$ has infinitely many connected components denoted by $(C_n)_{n \geq 1}$. For any $n \geq 1$, let $e_n \in E(\overline{C_n}) \cap Y$ and $a_n \in E(\overline{C_n}) \setminus \{e_n\} \subset E(X)$. For any $n \geq 1$, $(e_n, a_n] \subset C_n$ so $((e_n, a_n])_{n \geq 1}$ is a pairwise disjoint connected subsets in X . By Lemma 2.5, $\lim_{n \rightarrow +\infty} \text{diam}((e_n, a_n]) = 0$ hence $\lim_{n \rightarrow +\infty} d(a_n, e_n) = 0$. Since $E(X)$ is closed, we may assume that the sequence $(a_n)_{n \geq 1}$ converges to a point $a \in E(X)'$ hence $(e_n)_{n \geq 1}$ converges to $a \in E(Y)$. Since $E(Y)$ is closed then $a \in E(Y) \cap E(X)'$, absurd. Let prove that Y is a tree. By Lemma 2.10, we have $E(Y)' \subset E(X)' \cap E(Y)$ then $E(Y)' = \emptyset$, since $E(Y)$ is closed then $E(Y)$ is finite. Hence Y is a tree. Now, denote by $C_1, C_2, \dots, C_k$; $k > 0$ the connected components of $X \setminus Y$. For any $1 \leq n \leq k$; denote by $\{e_n\} = Y \cap \overline{C_n}$. For any $n, m \in \{1, 2, \dots, k\}$, $n \sim m \Leftrightarrow e_n = e_m$. For any $1 \leq n \leq k$, let $D_n := \bigcup_{m \sim n} \overline{C_m}$. Then we obtain a pairwise disjoint subdendrites $D_1, D_2, \dots, D_s$; $s \geq 1$ satisfying the conditions of the Lemma. $\square$

Proof of Proposition 5.2. We distinguish two cases.

Case 1: $M \cap \text{Fix}(f) \neq \emptyset$.

Let $a \in \text{Fix}(f) \cap M$. Denote by $F_a = \bigcup_{n=0}^{+\infty} f^{-n}(a) \cap M$ and $Y := [\overline{F_a}]$. We remark that by Theorem A, $a \in M \setminus L \subset M \setminus E(M)$, hence $2 \leq \text{ord}(a, M)$.

Lemma 5.6. *Let Assumption 5.1 be satisfied. Then $L \subset M \setminus Y$.*

Proof. **Claim 1.** For any $z \in F_a$, we have $[a, z] \cap L = \emptyset$.

Suppose contrary that $[a, z] \cap L \neq \emptyset$ for some $z \in f^{-p}(a) \cap M; p \geq 1$. Let y be such a point in the intersection above. Denote by $I = [a, y], J = [y, z]$ then we have $f^p(I) \cap f^p(J) \supset [a, f^p(y)]$. By Lemma 3.3, there is $i, j \geq 0$ such that $[a, f^{p+j}(y)] \supset [a, f^i(x)]$ so we have $f^{p+j}(I) \cap f^{p+j}(J) \supset [a, f^{i+r}(x)]$. Let $w \in L$ such that $z \in (y, w)$ then there is $r \geq 0$ such that $[a, z] \subset [a, f^{i+r}(x)]$. Then $f^{p+j+r}(I) \cap f^{p+j+r}(J) \supset [a, f^{i+r}(x)] \supset [a, z] = I \cup J$. Then I, J form an arc horseshoe for f , so $h(f) > 0$, absurd. This finish the proof Claim (1).

Claim 2. $L \cap \overline{F_a} = \emptyset$.

Suppose that there is $y \in L \cap \overline{F_a}$. First, we will prove that $f^n(y) \notin \overline{F_a}$ for some $n \geq 0$. Suppose contrary that $O_f(y) \subset \overline{F_a}$, then we have $\omega_f(y) \subset \overline{F_a}$, since $\omega_f(y) \subset L$ then it contain no periodic point, so $\omega_f(y)$ is uncountable. By Lemma 3.2, there is an open free arc I in X such that $\text{card}(\omega_f(y) \cap I) > 2$. Let z_1, z_2 two distinct points in $\omega_f(y) \cap I$ such that $z_1 \in (a, z_2)$ and I_1, I_2 two open disjoint arc in I containing z_1, z_2 , respectively. Then there is $k, p \geq 0$ such that $f^k(y) \in I_1$ and $f^p(y) \in I_2$. Since I_2 is open containing $f^p(y) \in \overline{F_a}$, then there is a point $z \in F_a \cap I_2$, hence $f^k(y) \in L \cap [a, z]$, this contradict Claim (1). Now, fix $n_0 \geq 1$ such that $f^{n_0}(y) \notin \overline{F_a}$. Since $X \setminus \overline{F_a}$ is a non empty open subset in X and since f^{n_0} is continuous, there is an open subset U of X containing y such that $f^{n_0}(U) \subset X \setminus \overline{F_a}$, let $t \in F_a \cap U$, then $f^{n_0}(t) \in f^{n_0}(U) \subset X \setminus \overline{F_a}$, but $f^{n_0}(t) \in F_a$, absurd. This finish the proof of Claim (2).

By Claims (1) and (2), we have for any $z \in \overline{F_a}$, $[a, z] \cap L = \emptyset$, so we have $L \subset M \setminus \bigcup_{z \in \overline{F_a}} [a, z]$, hence by Lemma 2.4 we obtain $L \subset M \setminus Y$. Since $E(Y) \subset \overline{F_a}$, $E(M)' \subset E(M) \subset L$ then $E(Y) \cap E(M)' = \emptyset$ then by Lemma 5.5, Y is a tree and $M \setminus Y$ has finitely many connected components. $\square$

Lemma 5.7. *Let Assumption 5.1 be satisfied. In (1) we suppose that $F_a \neq \{a\}$, then*

- (1) *for any $y \in L$, $(a, y] \cap F_a \neq \emptyset$,*
- (2) *$M \setminus F_a$ has finitely many connected components intersecting L , we denote it by $C_1, C_2, \dots, C_n$; $n \geq 2$ and for any $k \in \{1, 2, \dots, n\}$, we denote by $l_k = L \cap C_k$,*
- (3) *for any $1 \leq k \leq n$, there is a unique $j := \sigma(k) \in \{1, 2, \dots, n\}$ such that $f(C_k) \cap C_j \neq \emptyset$,*
- (4) *for any $1 \leq k \leq n$, $f(l_k) = l_{\sigma(k)}$,*
- (5) *for any $1 \leq k \leq n$, l_k is a clopen set relatively to L ,*
- (6) *σ is an n -cycle.*

Proof. (1) let $y \in L$ arbitrarily, $z \in F_a, z \neq a$ and $w \in L$ such that $z \in (a, w)$. By Lemma 3.3, there is $p, k \geq 0$ such that $[a, f^k(x)] \subset [a, f^p(y)] \subset f^p([a, y])$. Let $i \geq 0$ be such that $[a, z] \subset [a, f^{k+i}(x)]$ then $[a, z] \subset f^i([a, f^k(x)]) \subset$

$f^{p+i}([a, y])$ so $z \in f^{p+i}([a, y])$ hence there is $z_{-1} \in [a, y]$ such that $f^{p+i}(z_{-1}) = z \in F_a$ with $z_{-1} \neq a$. So $z_{-1} \in (a, y] \cap F_a$.

(2) If $F_a \neq \{a\}$. We have $F_a \subset Y$ so a connected subset of $M \setminus Y$ is also a connected subset of $M \setminus F_a$, since $L \subset M \setminus Y$ and by Lemma 5.5, $M \setminus Y$ has finitely many connected components so the connected subsets of $M \setminus F_a$ intersecting L are finite. If $F_a = \{a\}$, since $\text{ord}(a, M) < +\infty$ then $M \setminus \{a\}$ has finitely many connected components, (exactly $n := \text{ord}(a, M)$ components), and $L \subset M \setminus \{a\}$.

(3) Let $1 \leq k \leq n$, since $f(l_k) \subset f(C_k)$ and $f(l_k) \subset f(L) = \bigcup_{i=1}^n l_i$ so there is $1 \leq j \leq n$ such that $f(l_k) \cap l_j \neq \emptyset$ hence $f(C_k) \cap C_j \neq \emptyset$. Suppose that there is $1 \leq i \neq j \leq n$ such that $f(C_k) \cap C_i \neq \emptyset \neq f(C_k) \cap C_j$. Let $u \in f(C_k) \cap C_i, v \in f(C_k) \cap C_j$, then the subset $H := C_i \cup [u, v] \cup C_j$ is connected included in M since $[u, v] \subset M$. Since $f(C_k) \cap F_a = \emptyset$ for any $1 \leq k \leq n$ then $H \cap F_a = \emptyset$. By maximality of C_i and C_j we have $H \subset C_i$ and $H \subset C_j$ hence $H = C_i = C_j$, absurd.

(4) Let $1 \leq k \leq n$, for any $i \neq \sigma(k)$, $f(C_k) \cap C_i = \emptyset$ then $f(l_k) \cap l_i = \emptyset$, since $f(l_k) \subset L = \bigcup_{r=1}^n l_r$ hence $f(l_k) \subset l_{\sigma(k)}$. Suppose contrarily that $f(l_k) \subsetneq l_{\sigma(k)}$, since $f(L) = L$ then there is $i \neq k$ such that $f(l_i) \cap l_{\sigma(k)} \neq \emptyset$ hence so $\sigma(i) = \sigma(k)$. We obtain $f(L) = \bigcup_{r=1}^n f(l_r) \subset \bigcup_{r=1}^n l_{\sigma(r)} = \bigcup_{r=1; r \neq i}^n l_{\sigma(r)} \subsetneq L$, absurd. Then $f(l_k) = l_{\sigma(k)}$.

(5) Suppose that for some $1 \leq i \leq n$, $l_i \subsetneq \bar{l}_i$. Then there is $j \in \{1, 2, \dots, n\}; j \neq i$ such that $\bar{l}_i \cap l_j \neq \emptyset$ hence $\bar{C}_i \cap C_j \neq \emptyset$. Since C_i and C_j are disjoint then by Lemma 2.3 there is $z \in M$ such that $\bar{C}_i \cap C_j = \{z\}$. Since $C_i \subset (C_i \cup \{z\}) \subset \bar{C}_i$ then $C_i \cup \{z\}$ is connected, hence $C_i \cup \{z\} \cup C_j = C_i \cup C_j$ is connected disjoint with F_a , this contradict the maximality of C_i and C_j . Then $l_i = \bar{l}_i$ i. e. any l_i is closed in X hence in L . In the other hand, since $l_1, l_2, \dots, l_n$ are pairwise disjoint then $l_i = L \setminus \bigcup_{j=1; j \neq i}^n l_j$ is open relatively to L . We conclude that any l_i is a clopen set relatively to L .

(6) Let prove that σ is n -cycle. Suppose contrarily that for some $1 \leq s \leq n$, $\sigma^p(s) = s$ with $0 < p < n$. The subset $F := \bigcup_{k=0}^{p-1} l_{\sigma^k(s)}$ is proper and clopen relatively to L . We have $f(F) = \bigcup_{k=0}^{p-1} f(l_{\sigma^k(s)}) = \bigcup_{k=0}^{p-1} l_{\sigma^{k+1}(s)} = F$. Let $G = L \setminus F$ a non empty closed subset in L , $f(L \setminus G) \cap G = \overline{f(F)} \cap G = F \cap G = \emptyset$. This contradict Lemma 2.15. Hence $\forall s \in \{1, 2, \dots, n\}$, $O_\sigma(s) = \{1, 2, \dots, n\}$ then σ is an n -cycle. $\square$

Lemma 5.8. Denote by $s_1 := [l_1] \subset C_1$. The subset $S := \bigcup_{k=0}^{+\infty} f^{kn}(s_1)$ satisfies conditions (i) – (iv) of the Proposition 5.2.

Proof. First, S is connected. Indeed, since s_1 is a subdendrite of X then it is so for $f^{kn}(s_1), \forall k \geq 0$. We have $f^n(s_1) \supset f^n(l_1) = l_{\sigma^n(1)} = l_1$ hence

$f^n(s_1) \supset s_1$, so $(f^{kn}(s_1))_{k \geq 0}$ is an increasing sequence of connected subsets then S is connected.

(i) $S, f(S), \dots, f^{n-1}(S)$ are pairwise disjoint. Indeed, since $f^{-1}(F_a) = F_a$ and $s_1 \cap F_a = \emptyset$ then $f^i(s_1) \cap F_a = \emptyset, \forall i \geq 0$ hence $f^k(S) \cap F_a = \emptyset; \forall 0 \leq k \leq n-1$. Suppose that $f^i(S) \cap f^j(S) \neq \emptyset$ for some $0 \leq i \neq j \leq n-1$ then $f^i(S) \cup f^j(S)$ is connected. Let $u \in l_{\sigma^i(1)}, v \in l_{\sigma^j(1)}$, since $f^k(S) \supset l_{\sigma^k(1)}$ for any $k \geq 0$ then $[u, v] \subset (f^i(S) \cup f^j(S)) \cap M$. The subset $K := C_{\sigma^i(1)} \cup [u, v] \cup C_{\sigma^j(1)}$ is connected in M disjoint with F_a , by maximality of $C_{\sigma^i(1)}$ and $C_{\sigma^j(1)}$ we obtain $K \subset C_{\sigma^i(1)}$ and $K \subset C_{\sigma^j(1)}$ hence $C_{\sigma^i(1)} = C_{\sigma^j(1)}$, absurd.

(ii) $f^n(S) = \bigcup_{k=0}^{+\infty} f^{(k+1)n}(s_1) = \bigcup_{k=0}^{+\infty} f^{nk}(s_1) = S$.

(iii) Since $l_1 \subset S$ then $L = \bigcup_{k=0}^{n-1} l_{\sigma^k(1)} \subset \bigcup_{k=0}^{n-1} f^k(S)$.

(iv) $L \cap f^k(S) = l_{\sigma^k(1)}$ for any $0 \leq k \leq n-1$. Indeed, we have $f^k(S) \supset f^k(l_k) = l_{\sigma^k(1)}; \forall k \geq 0$ and since $S, f(S), \dots, f^{n-1}(S)$ are pairwise disjoint then $L \cap f^k(S) = l_{\sigma^k(1)}$, so $f(L \cap f^k(S)) = f(l_{\sigma^k(1)}) = l_{\sigma^{k+1}(1)} = L \cap f^{k+1}(S)$. $\square$

This finish the proof of Case 1.

Case 2: $M \cap \text{Fix}(f) = \emptyset$.

In the following Lemma we will use the notations from [18].

Lemma 5.9. *If M does not contains a fixed point then there is a fixed point $z \in X \setminus M$ such that $M \subset \psi_f^{-1}(z)$.*

Proof. Let $y \in M$ such that $\psi_f(y) = z \in \text{Fix}(f)$. Suppose contrarily that $M \not\subset \psi_f^{-1}(z)$ then $M \cap (X \setminus \psi_f^{-1}(z)) \neq \emptyset$ since $M \cap \psi_f^{-1}(z) \neq \emptyset$ and M is connected then $M \cap \partial(\psi_f^{-1}(z)) \neq \emptyset$ but $\partial(\psi_f^{-1}(z)) \subset \text{Fix}(f)$ hence $M \cap \text{Fix}(f) \neq \emptyset$, absurd. Then $M \subset \psi_f^{-1}(z)$. It follow that $\forall y \in M, \psi_f(y) = z$. This finish the proof of the Lemma. $\square$

Now, let $z_0 := r_M(z)$ where r_M is the retraction map (See [19]) and denote by $F_{z_0} := \bigcup_{n=0}^{+\infty} f^{-n}(z_0) \cap M$ and $Y_0 := \overline{F_{z_0}}$. We remark that $F_{z_0} \cap L = \emptyset$ (so $\text{ord}(z_0, M) \geq 2$), F_{z_0} is non empty and any arc joining a point in M and $f(z_0)$ contains z_0 .

We have an analogous results compared to Case 1.

Lemma 5.10. *Let Assumption 5.1 be satisfied. Then we have the following properties:*

- (1) $L \subset M \setminus Y_0$,
- (2) $M \setminus Y_0$ has finitely many connected components. The connected components of $M \setminus F_{z_0}$ intersecting L are denoted by $C_1, C_2, \dots, C_n$ where $n > 1$, we denote by $l_k = L \cap C_k$,
- (3) for any $1 \leq k \leq n$, there is a unique $j := \sigma(k) \in \{1, 2, \dots, n\}$ such that $f(C_k) \cap C_j \neq \emptyset$,
- (4) for any $1 \leq k \leq n$, $f(l_k) = l_{\sigma(k)}$,
- (5) for any $1 \leq k \leq n$, l_k is a clopen set relatively to L ,

- (6) σ is an n -cycle,
- (7) Let $S = \bigcup_{k=0}^{+\infty} f^{kn}([l_1])$. Then
 - (a) $S, f(S), \dots, f^{n-1}(S)$ are a pairwise disjoint connected subsets,
 - (b) $f^n(S) = S$,
 - (c) $L \subset \bigcup_{i=0}^{n-1} f^i(S)$

Proof. (1) **Claim 1.** There is $n_0 \geq 0$ such that $\forall n \geq n_0$ we have $z_0 \in (f(z_0), f^n(x))$.

Since $z_0 \notin L$ and L is compact then $\delta := d(z_0, L) > 0$. Remark that for any $y \in L, z_0 \in (y, f(z_0))$. Then $\forall y \in L, \exists 0 < \mu = \mu(y) < \delta$ such that $\forall t \in B(y, \mu), z_0 \in (t, f(z_0))$. The subset $V := \bigcup_{y \in L} B(y, \mu(y))$ is an open neighborhood of L disjoint with z_0 . There is $n_0 \geq 0$ such that $\forall n \geq n_0, f^n(x) \in V$. Then $\forall n \geq n_0$, there is $y \in L, f^n(x) \in B(y, \mu(y))$ hence $z_0 \in (f^n(x), f(z_0))$.

Claim 2. For any $t \in F_{z_0}, [z_0, t] \cap L = \emptyset$.

Suppose contrarily that there is $t \in f^{-k}(z_0) \cap M; k > 0$ and $y \in L$ such that $y \in [z_0, t]$. Denote by $I = [z_0, y], J = [y, t]$. We have $f^k(J) \supset [z_0, f^k(y)]$ then $f^{k+1}(J) \supset [f(z_0), f^{k+1}(y)] \ni z_0$ hence $f^{k+1}(J) \supset [z_0, f^{k+1}(y)]$ so by induction, for any $n \geq k, f^n(J) \supset [z_0, f^n(y)]$. Similarly, we have $f(I) \supset [f(z_0), f(y)] \ni z_0$ then $f(I) \supset [z_0, f(y)]$. Hence we prove by induction that for any $n \geq 0, f^n(I) \supset [z_0, f^n(y)]$. As the proof of Lemma 5.6, Claim 1, we show easily that I, J form an arc horseshoe hence $h(f) > 0$, a contradiction.

Claim 3. $\overline{F_{z_0}} \cap L = \emptyset$.

Suppose contrarily that there is $y \in L \cap \overline{F_{z_0}}$. Let $(z_i)_{i>0}$ be a sequence in F_{z_0} converging to y and $f^{n_i}(z_i) = z_0; \forall i > 0$. Let prove that there is a neighborhood V of y in X and $k \geq 0$ such that $f^k(V) \subset I$ where I is a free open arc in X such that $[z_0, s] \cap L \neq \emptyset$ for some $s \in F_{z_0}$. Indeed, since $\omega_f(y)$ is uncountable then there is a free open arc I in X such that $I \subset M$ and $\omega_f(y) \cap I$ is uncountable. Let y_1, y_2 two distincts points in the intersection such that $y_1 \in (z_0, y_2)$ and I_1, I_2 two open disjoint arc in I containing y_1, y_2 respectively. Let $k, p > 0$ such that $f^k(y) \in I_1, f^p(y) \in I_2$. By continuity of f^p there is an open set, V , in X containing y such that $f^p(V) \subset I_2$. If the sequence $(n_i)_{i>0}$ is bounded then by taking a subsequence we may assume that $n_i =: m$ for any $i > 0$. So we have $f^m(z_i) = z_0, \forall i > 0$, since $z_i \rightarrow_{i \rightarrow +\infty} y$ then $f^m(y) = z_0 \in L$, absurd. If the sequence $(n_i)_{i>0}$ is unbounded, we may assume that it is non decreasing and $n_i > p, \forall i > 0$. Let $i > 0$ such that $z_i \in V$ then $f^p(z_i) \in f^p(V) \subset I_2 \subset M$. So we have $f^k(y) \in (z_0, f^p(z_i))$ and $f^{n_i-p}(f^p(z_i)) = z_0$ then $f^p(z_i) \in F_{z_0}$, this contradict Claim 2. Hence we finish the proof of Claim 3.

Now, we have $L \subset M \setminus [z_0, t], \forall t \in \overline{F_{z_0}}$ then $L \subset M \setminus \bigcup_{t \in \overline{F_{z_0}}} [z_0, t] = M \setminus Y_0$.

(2) Y_0 is a subdendrite of M satisfying $E(Y_0) \cap E(M)' \subset \overline{F_{z_0}} \cap L = \emptyset$ then $M \setminus Y_0$ has finitely many connected components.

The proof of (3) – (7) are similar as in Lemmas 5.7 and 5.8.

□

5.1. Proof of Theorem B. Let S be a connected subset of X satisfying conditions (i) – (iii) of proposition 5.2. Denote by $\alpha_1 := n_1$ and $D_1 = \overline{S}$. Then we have $L \subset \cup_{i=0}^{\alpha_1-1} f^i(D_1)$. Set $f_1 := f|_{D_1}^{\alpha_1} : D_1 \rightarrow D_1$, it is a dendrite map such that $E(D_1)$ is countable closed and $E(D_1)'$ finite. Let $t_1 = f^{p_1}(x) \in D_1$. Since $\omega_{f_1}(t_1)$ is uncountable then by Proposition 5.2 there is a pairwise disjoint connected subsets $S_2, f_1(S_2), \dots, f_1^{n_2-1}(S_2); n_2 \geq 2$, $f_1^{n_2}(S_2) = S_2$ in D_1 such that $\omega_{f_1}(t_1) \subset \cup_{i=0}^{n_2-1} f_1^i(S_2)$. So for any $0 \leq j \leq n_1 - 1$, $\omega_{f^{n_1}}(f^j(t_1)) \subset \cup_{i=0}^{n_2-1} f^{in_1+j}(S_2) \subset \cup_{i=0}^{n_2-1} f^{in_1+j}(D_2)$ where $D_2 = \overline{S_2}$. Hence $L = \omega_f(t_1) \subset \cup_{j=0}^{n_1-1} \cup_{i=0}^{n_2-1} f^{in_1+j}(D_2) = \cup_{i=0}^{\alpha_2-1} f^i(D_2)$. The subdendrite D_2 has period $\alpha_2 := n_1 n_2$ and $\cup_{k=0}^{n_2-1} f^{i\alpha_1}(D_2) \subset D_1$. Since $\text{card}(f^i(D_1) \cap f^j(D_1)) \leq 1$ for any $0 \leq i \neq j \leq \alpha_1 - 1$ then $\text{card}(f^i(D_2) \cap f^j(D_2)) \leq 1$ for any $0 \leq i \neq j \leq \alpha_2 - 1$. By induction we build a sequence of subdendrites $(D_k)_{k \geq 1}$ satisfying the conditions of Theorem B.

6. On Li-Yorke pair implies chaos.

To prove Theorem C we need the following Lemma.

Lemma 6.1. *We use the notations from Theorem B. There is $k > 0$ and $0 \leq i < \alpha_k$ such that $f^i(D_k)$ is a free arc.*

Proof. Case 1. If X is a tree. The number $N := \sum_{b \in B(X)} \text{ord}(b, X)$ is finite, let $k > 0$ such that $\alpha_k > N$. Then there is $0 \leq i < \alpha_k$ such that $f^i(D_k) \cap B(X) = \emptyset$ then $f^i(D_k)$ is a free arc in X .

Case 2. If X is not a tree. Let $d := \text{card}(E(X)') \geq 1$. For any $k \geq 1$ we have $n_k \geq 2$ so $\alpha_k \geq 2^k$. Let $k > 0$ such that $d < 2^k$. Since $\text{card}(f^i(D_k) \cap f^j(D_k)) \leq 1$ then there is $0 \leq i < \alpha_k$ such that $f^i(D_k) \cap E(X)' = \emptyset$. By Lemma 5.5, $f^i(D_k)$ is a tree. By Case 1, there is $p \geq k$ and $0 \leq j < \alpha_p$ such that $f^j(D_p)$ is an arc included in $f^i(D_k)$. Since $f^j(D_p) \cap E(X)' = \emptyset$ then the set $f^i(D_p) \cap B(X)$ is finite. So there is $q > p$ and $0 \leq s < \alpha_q$ such that $f^s(D_q)$ is a free arc in X . $\square$

Proof of Theorem C. Let (x, y) be a Li Yorke pair for f such that $L := \omega_f(x)$ is uncountable. Denote by $I = f^i(D_k) = [u, v]$ a free arc in X . There is $n \geq 0$ such that $x_n \in I$. Denote by $d = \alpha_k$, then $g = f|_I^d : I \rightarrow I$ is an interval map. Since (x, y) is a Li Yorke pair for f then (x_n, y_n) is proximal for f^d . So $\omega_{f^d}(x_n) \cap \omega_{f^d}(y_n) \neq \emptyset$. We distinguish two cases:

Case 1: $O_{f^d}(y_n) \cap I \neq \emptyset$. Let $k \geq 0$ such that $f^{dk}(y_n) \in I$. So $(f^{dk}(x_n), f^{dk}(y_n)) \in I^2$ is a Li Yorke pair for $g : I \rightarrow I$, by [17] g is chaotic hence f is chaotic.

Case 2. $O_{fd}(y_n) \cap I = \emptyset$. Since $\omega_{fd}(x_n) \cap \omega_{fd}(y_n) \neq \emptyset$ and $\omega_{fd}(x_n) \subset I$ then $\omega_{fd}(y_n) \cap I \subset \{u, v\}$ so $\omega_{fd}(x_n) \cap \omega_{fd}(y_n)$ is finite and f^d -invariant so $\omega_{fd}(x_n)$ contains a periodic point, say u . Then $(x_n, u) \in I^2$ is a Li Yorke pair for g , by [17] g is chaotic, hence f is chaotic.

7. Example

Example 4. (Due to I. Naghmouchi) We will construct a dendrite X with $E(X)$ is countable closed set such that $(E(X))^{(2)}$ is reduced to one point and a map f on X having a Li-Yorke pair but not Li-Yorke chaotic.

Proof. **Construction of the dendrite X .**

Denote by $D_0 = ([-1, 1] \times \{0\}) \cup (\bigcup_{n \geq 1} \{1 - \frac{1}{n}\} \times [0, \frac{1}{n}])$. Let $I = [(0, 0), (\frac{1}{2}, -1)]$ and $X = I \cup (\bigcup_{n \geq 1} X_n)$ where for any $n \geq 1$,

- X_n is a homeomorphic copy of D_0 , by a homeomorphism φ_n ,
- $a_k^n := \varphi_n((1 - \frac{1}{k}, \frac{1}{k}))$, $\forall k \geq 1$ and $a^n = \varphi_n((1, 0))$, $a^0 = (0, 0)$,
- $X_n \cap I = \{b_n\}$ where $\varphi_n((-1, 0)) = b_n$,
- the sequence $(X_n)_{n \geq 1}$ are pairwise disjoint and $\lim_{n \rightarrow +\infty} \text{diam}(X_n) = 0$,
- $c_k^n = \varphi_n((1 - \frac{1}{k}, 0))$, $\forall n, k \geq 1$.

We can see that the set X is a dendrite such that $E(X) = \{a_k^n; n, k \geq 1\} \cup \{a^n; n \geq 1\} \cup \{a^0\}$, $E(X)' = \{a^n; n \geq 1\} \cup \{a^0\}$ and $E(X)^{(2)} = \{a^0\}$.


Construction of the map f . (See Figure 3)

The map f is defined as follows:

- $f(a^1) = f(a^0) = a^0$ and $f(b_1) = b_1$,
- $f([b_2, c_1^1]) = \{b_1\}$,
- $\forall n \geq 1, f([b_n, b_{n+1}]) = \{b_n\}$,
- $\forall n, k \geq 1, f$ sent linearly $[b_{n+1}, a^{n+1}]$ to $[b_n, a^n]$ such that $f(c_k^{n+1}) = c_{k+1}^n$ and f sent linearly $[c_k^{n+1}, a_k^{n+1}]$ to $[c_{k+1}^n, a_{k+1}^n]$,
- $\forall k \geq 1, f$ sent linearly $[c_1^1, a^1]$ to $[b_1, a^0]$ such that $f(c_k^1) = b_k$ and f sent linearly $[c_k^1, a_k^1]$ to $[b_k, a_1^{k+1}]$.

We see that f is a dendrite map having a Li-Yorke pair (take for example (a^0, a_1^1)) but f is not Li-Yorke chaotic. Indeed, if $(x, y) \in X \times X$ is a Li yorke pair then $(x, y) \in E(X)^2$.

□

FIGURE 3. Dendrite with $E(X)'$ is infinite

Lemma 7.1. *Every dendrite X with closed set of endpoints and $E(X)^{(2)}$ is non empty contains a subdendrite Y with closed set of endpoints and $E(Y)^{(2)}$ is reduced to one point.*

Proof. Claim. If F is a closed set such that $F^{(2)}$ is non empty then it contain a closed subset A such that $A^{(2)}$ is reduced to one point.

Let $e \in F^{(2)}$, there is a sequence of pairwise different elements $(e_n)_{n \geq 0}$ in F' converging to e . Let $(\varepsilon_n)_{n \geq 0}$ a sequence of non negative reels such that the sequence of balls $(B(e_n, \varepsilon_n))_{n \geq 0}$ are pairwise disjoint. For any $n \geq 0$, since $e_n \in F'$ then there is a sequence $(e_n^k)_{k \geq 0}$ in $B(e_n, \varepsilon_n) \cap F$ converging to e_n . Hence the set $A = \{e_n^k; n, k \geq 0\} \cup \{e_n; n \geq 0\} \cup \{e\}$ satisfy the condition of the Claim. Now, let A be a closed subset in $E(X)$ such that $A^{(2)}$ is one point. By Lemma 2.4, $Y := [A]$ is a subdendrite of X with $E(Y) = A$. $\square$

By Lemma 7.1, there is a non chaotic dendrite map $g : Y \rightarrow Y$ having a Li-Yorke pair $(x, y) \in Y^2$. Let $r_Y : X \rightarrow Y$ be the retraction map and $f = g \circ r_Y : X \rightarrow Y \subset X$. Then (x, y) is also a Li Yorke pair for f but f is not Li-Yorke chaotic. $\square$

REFERENCES

- [1] R. L. Adler, A. G. Konheim, M. H. McAndrew, *Topological entropy*, Trans. Amer. Math. Soc. **114**, (1965), 309–319.
- [2] L. Alseda, S. Kolyada, J. Libre, L. Snoha, *Entropy and periodic points for transitive maps*, Trans. Amer. Math. Soc. **351** (1999), 1551–1573.

- [3] D. Arévalo, W. J. Charatonik, P. P. Covarrubias, L. Simn, *Dendrites with a closed set of endpoints*, Top. App. **115** (2001), 1–17.
- [4] F. Blanchard, E. Glasner, S. Kolyada, A. Maass, *On Li-Yorke pairs*, J. Reine Angew. Math. **547** (2002), 51–68.
- [5] F. Blanchard, W. Huang and L. Snoha, *Topological size of scrambled sets*, Colloq. Math. **110** (2008), 293–361.
- [6] L. S. Block, W. A. Coppel, *Dynamics in One Dimension*, Lecture Notes in Math, 1513. Springer-Verlag, 1992.
- [7] R. Bowen, *Entropy for group endomorphisms and homogeneous spaces*, Trans. Amer. Math. Soc. **153** (1971), 401–414.
- [8] J. J. Charatonik, W. J. Charatonik, J. R. Prajs, *Mapping hierarchy for dendrites*, Dissertationes Math. (Rozprawy Mat.) **333** (1994), 52 pp.
- [9] W. J. Charatonik, E. P. Wright, S. S. Zarafidou, *Dendrites with a countable set of endpoints and universality*, Houston J. of Math, **39** (2013), 651–666.
- [10] E. I. Dinaburg, *The relation between topological entropy and metric entropy*, Doklady Akademii Nauk SSSR **190** (1970), 19–22.
- [11] G. L. Forti, L. Paganoni, J. Smital, *Strange triangular maps of the square*, Bull. Austral. Math. Soc. **51** (1995), 395–415.
- [12] X. C. Fu, Z. M. Wang, *The construction of chaotic subshifts*, J. Nonlin. Dyn. Sci. Technol. **4**(3) (1997) 127–132 (in Chinese).
- [13] J. L. G. Guirao, M. Lampart, *Li and Yorke chaos with respect to the cardinality of the scrambled sets*, Chaos Solitons Fractals **24** (2005), 1203–1206.
- [14] R. Hric, M. Malek, *Omega limit sets and distributional chaos on graphs*, Top. App. **153** (2006), 2469–2475.
- [15] Z. Kocan, *Chaos on one-dimensional compact metric spaces*, Int. J. Bifurcation and Chaos, **22** (2012), 125–259.
- [16] Z. Kocan, V. K. Kurkova, M. Malek, *Entropy, horseshoes and homoclinic trajectories on trees, graphs and dendrites*, Ergodic Theory Dynam. Systems, **31** (2011), 165–175.
- [17] M. Kuchta, J. Smital, *Two-point scrambled set implies chaos*. European Conference on Iteration Theory (Caldes de Malavella, 1987), p. 427–430. World Sci. Publishing, Teaneck, NJ, 1989.
- [18] J. Mai, E. Shi, $\overline{R} = \overline{P}$ for maps of dendrites X with $\text{Card}E(X) < c$, Int. J. Bifurcation and Chaos, **19** (2009), 1391–1396.
- [19] S. B. Nadler, *Continuum Theory: An Introduction*, Monogr. Textb. Pure Appl. Math., vol. 158, Marcel Dekker, Inc., New York, 1992.
- [20] J. Nikiel, *A characterisation of dendroids with uncountable many endpoints in the classical sense*, Houston J. Math. **9** (1983), 421–432.
- [21] S. Ruelle, L. Snoha, *For graph maps, one scrambled pair implies Li-Yorke chaos*, Proc. Amer. Math. Soc. **142** (2014), 2087–2100.
- [22] A. N. Sarkovskii, *The behavior of a map in a neighborhood of an attracting set*, Ukrain. Mat. Z. **18** (1966), 60–83. (Russian).
- [23] J. Smital, *Chaotic functions with zero topological entropy*, Trans. Amer. Math. Soc. **297** (1986), 269–282.

GHASSEN ASKRI, UNIVERSITY OF CARTHAGE, FACULTY OF SCIENCES OF BIZERTE,
DEPARTMENT OF MATHEMATICS, JARZOUNA, 7021, TUNISIA
E-mail address: askri.ghassen@hotmail.fr